

Stefanie K. Taillon
Acting Secretary of Natural and Historic Resources

Matthew S. Wells
Director

Andrew W. Smith
Chief Deputy Director



Frank N. Stovall
*Deputy Director
for Operations*

Darryl Glover
*Deputy Director for
Dam Safety,
Floodplain Management and
Soil and Water Conservation*

Laura Ellis
*Deputy Director for
Administration and Finance*

COMMONWEALTH of VIRGINIA
DEPARTMENT OF CONSERVATION AND RECREATION

Approved February 13, 2025

Date Received by DCR February 11, 2025

Randy Johnson
GCC Locust Grove
1000 Germanna Point Drive
Fredericksburg, Va 22408

Your nutrient management plan (NMP) dated March 1, 2025 located in Locust Grove has been approved by the Virginia Department of Conservation and Recreation (DCR). The approved plan is for 8.83 acres. Only nutrient recommendations for applications to be made after the date of this letter are approved by this letter. Your NMP was written by Sara Shelton, a nutrient management planner certified by DCR.

This site has not been inspected by DCR and this approval is contingent upon site conditions being as stated in the NMP. Any revisions to this plan must be approved by DCR. Any change in personnel resulting in a change to the plan manager should be reported to the Certified Nutrient Management Planner who will then make DCR aware. Please note that this letter should be kept with the NMP and supporting documentation including nutrient application records. This plan expires on March 1, 2028. Please feel free to contact me with any questions or concerns regarding this approval.

Best regards,

James Janney

James Janney
Urban Nutrient Management Coordinator
Division of Soil and Water Conservation
600 East Main Street, 24th Floor
Richmond VA 23219
(804) 513-5958

Nutrient Management Plan for the Locust Grove Campus Germanna Community College

Prepared for:

**Randy Johnson
Director of Facilities
Germanna Community College
1000 Germanna Point Drive
Fredericksburg, VA 22408
540.423.9046**

Prepared By:

Sara J. Shelton/Certified Nutrient Management Planner - Certification No. 943

**Wetland Studies and Solutions, Inc.
1620 Brook Road
Richmond, Virginia 23220**

Location Information	
Physical Address	<i>2130 Germanna Highway</i>
City State Zip	<i>Locust Grove, VA 22508</i>
Coordinates	<i>38° N 22' 30.53"</i>
NAD 83 Deg Min Sec	<i>77° W 46' 52.91"</i>
VAHU6 Watershed Code	<i>RA-42 – Rapidan River-Fields Run</i>
County	<i>Orange</i>

Square Footage of Management Areas	
Total	<i>8.83 acres (384,823.52 ft²)</i>
Area 1	<i>5.40 acres (235,525.00 ft²)</i>
Area 1	<i>3.43 acres (149,298.52 ft²)</i>

Plan Start Date	<i>March 1, 2025</i>
Plan End Date	<i>March 1, 2028</i>
Planner Signature	<i>Sara Shelton</i>

TABLE OF CONTENTS

1.0	INTRODUCTION AND SITE DESCRIPTION	1
1.1	Introduction.....	1
1.2	Site Description	1
1.3	Current and Future Turf Maintenance	1
2.0	SOIL SAMPLING AND ANALYSIS	1
3.0	NUTRIENT MANAGEMENT AREAS	2
3.1	Nutrient and Liming Applications.....	2
3.1.1	Nitrogen, Phosphorous and Potassium	2
3.1.2	Lime and pH.....	3
3.2	Problem Turfgrass Areas and Temporarily Inactive Nutrient Management Areas.....	3
3.3	Selection of Fertilizers	3
3.4	Pre- and Post-Emergent Herbicides.....	4
3.5	Precautions for Fertilizer Applications	4
4.0	ENVIRONMENTALLY SENSITIVE AREAS AND RECOMMENDED BUFFERS	4
5.0	OTHER TURF MANAGEMENT CONSIDERATIONS	5
6.0	RECORDKEEPING.....	5
7.0	REFERENCES.....	6

Tables

Table 1: Soil Test Summaries

Table 2: Nutrient Application Worksheet (Schedule 1)

Table 3: Nutrient Application Worksheet (Schedule 2)

Figures

Figure 1: Project Location

Figure 2: Soil Sampling Areas

Figure 3: Nutrient Management Areas

Figure 4: Liming Requirements

Appendices

Appendix A: Laboratory Soil Test Results

Appendix B: Application Record Forms

1.0 INTRODUCTION AND SITE DESCRIPTION

1.1 Introduction

This Nutrient Management Plan (NMP) is for the Locust Grove Campus of the Germanna Community College (GCC) located on Route 3 in between Culpeper and Locust Grove in Orange County, Virginia (Figure 1). The entire property is 99.60 acres.

This NMP addresses only the nutrient management of turfgrass. Management of other vegetated areas containing trees, flowering ornamentals, small shrubs and groundcovers, is performed by each facility based on very site-specific conditions including but not limited to the type and status of vegetated areas, annual soil testing, and the occurrence of pests and weeds. This NMP is effective until March 1, 2028, or until major renovation or other changes to maintenance practices occur. This NMP should be used as a resource for planning the quantity and timing of turfgrass nutrient application based on sound agronomic practices.

1.2 Site Description

The Campus contains turfgrass in many areas including around campus buildings, along roadways and around and within parking lots. The grounds are managed at a moderate maintenance level. Turf areas consist of blends of fescue including the mixed-use athletic field.

1.3 Current and Future Turf Maintenance

Turf maintenance is performed by GCC staff and other outside contractors when needed including mowing, herbicide, fertilizer and lime applications, as well as aeration and overseeding. It will be the responsibility of the Director of Facilities to ensure the management plan is followed.

2.0 SOIL SAMPLING AND ANALYSIS

Although most of the soils in the turf areas have been modified by cut and fill activities, the soils still retain most of the characteristics presented in the U.S. Department of Agriculture (USDA) soil survey and may still be classified as clay loams.

Soil samples were collected on December 6, 2024, from three (3) different turfgrass areas across the campus and submitted for laboratory analysis including pH, buffer pH, phosphorus and potassium, and other soil properties. Figure 2 shows the locations of the soil sampling areas as well as environmentally sensitive areas and Table 1 presents the laboratory results. Appendix A presents the soil laboratory data. No sampling was performed within wooded or landscaped areas.

Soil laboratory results were converted into nutrient management ratings based on the Virginia Nutrient Management Standards and Criteria (VNMS&C). Soil phosphorous levels were all L- and potassium concentrations ranged from L to M. Soil pH ranged from 5.3 to 5.7 Standard Units (SU), with all below the target level pH of 6.2.

3.0 NUTRIENT MANAGEMENT AREAS

Based on the soil test results, current turf conditions, the intensity of use, and overall visibility and aesthetic considerations, two (2) Nutrient Management Areas (NMA) at the Locust Grove Campus have been established for this NMP. The number of nutrient management areas will facilitate effective management, protect water quality, and maintain healthy turf. Figure 3 shows the nutrient management areas and Figure 4 shows the liming areas. Table 2 presents the application schedule for the nutrient management and liming areas, discussed in greater detail in Section 3.1 below.

3.1 Nutrient and Liming Applications

3.1.1 Nitrogen, Phosphorous and Potassium

Nitrogen, phosphorous (P₂O₅) and potassium (K₂O) are the three macronutrients essential for healthy turf and, along with lime applications, are the central focus of the NMP. Phosphorous and potassium recommendations are based on the soil laboratory results. Nitrogen recommendations are based on the turfgrass needs, not soil test results, which vary based on the type of turfgrass (cool vs. warm season) and level of management (standard vs. intensive). Recommended rates and timing of all three macronutrients are based on the VNMS&C. This NMP uses most restrictive application rate for each NMA based on individual sample results where multiple sampling areas are part of the same NMA.

The acceptable window for nitrogen application for cool season fescue turf at the Locust Grove Campus is six weeks prior to the last spring average frost and six weeks after the first fall average frost from March 4th until December 8th. Although aggressive spring and summer nitrogen fertilization can result in lush, dark green foliage, this occurs at the expense of the turf root system. Turf with an inadequate root system will then struggle in the summer heat and moisture conditions. Additionally, too much nitrogen in spring and summer for cool season turf can result in leaching or runoff to nearby waterbodies. The bulk of nitrogen should be applied in monthly increments from September through November.

As phosphorous and potassium are not as mobile as nitrogen and generally reside in soil for longer periods of time, the application timing of these two macronutrients is not as critical as nitrogen. Incremental applications of these nutrients from September to November are recommended.

3.1.2 Lime and pH

Soil acidity is critical to plants because it affects the availability of nutrients in the soil and potential leaching of nutrients from the soil. Cool season fescue prefers a soil pH that is slightly acidic, at a level of approximately 6.2 Standard Units (SU). Periodic lime applications are necessary for many Virginia soils to correct low pH, add buffering capacity, to provide secondary nutrients calcium and magnesium as well as some micronutrients. Liming rates are based on the soil test pH and the buffer indices.

According to the laboratory results, lime applications are not needed for the first year of this NMP. Liming recommendations are only for the first year following sampling. The soil should be tested for soil pH and Buffer pH in the late fall to winter each following year to determine if liming is necessary following the initial recommended liming.

3.2 Problem Turfgrass Areas and Temporarily Inactive Nutrient Management Areas

Turf in several areas across the campus appears thin and appears to need additions of topsoil and overseeding. Areas of ineffective groundcover and all active construction sites should be temporarily removed from active nutrient management until corrective measures can be applied to improve the turfgrass or groundcover conditions or the construction site is returned to active management. At the time of the development of this NMP, there were no active construction sites or turf areas that were designated as ineffective and removed from active management.

Corrective action options vary by area but may include additional soil amendments (compost/topsoil), aeration or shallow tilling, and the use of mulch, turf mats and blankets. Alternative landscaping such as groundcovers, pavers, and other hardscape treatments may be the best alternative for some areas. If turfgrass is the desired vegetative cover, the soil should be retested for soil and buffer pH and adjusted accordingly with limestone as part of corrective action.

3.3 Selection of Fertilizers

Specific fertilizers have not been selected as a part of this NMP to provide greater flexibility and cost savings. The landscape contractor has the option to select either commonly used fertilizer blends that they may already have in stock or are readily available, or they can use custom blends, a common practice in the commercial landscaping industry. Slow-release nitrogen containing fertilizers are recommended. This NMP will require revision should the Director of Facilities decide to use animal manures or Class B biosolids (not of exceptional quality).

Provided the maximum rate of nitrogen per application and the total annual rates of all three nutrients are not exceeded as detailed in Table 2, the Facility staff and/or a landscape contractor may use their discretion with the exact ratio of nutrients applied per application.

3.4 Pre- and Post-Emergent Herbicides

Weed control is a necessary requirement for healthy turf. Herbicides with nitrogen included may be used in the spring provided the application of nitrogen follows the amount allowed by this NMP and the VNMS&C. However, additional straight application of herbicides without nitrogen additives may be required.

3.5 Precautions for Fertilizer Applications

General precautions for fertilizer application include:

- Avoid applying fertilizers on steep slopes 48-hours prior to a rain event.
- Do not apply fertilizers to frozen or snow-covered ground, nor should they ever be used as ice melt.
- Avoid/minimize application of fertilizers to impervious areas such as parking lots, roads, and sidewalks, and within 25 feet of environmentally sensitive areas and stormwater collection/management facilities.
- Remove any granular materials that land on impervious surfaces by sweeping and collecting, and either put the collected material back in the bag or spread it onto the turf.

4.0 ENVIRONMENTALLY SENSITIVE AREAS AND RECOMMENDED BUFFERS

A small unnamed tributary to the Rapidan River was identified as an environmentally sensitive area on the Locust Grove Campus as shown on Figures 2 through 4. A no-fertilizer/pesticide application buffer area of at least 25 feet and preferably 50 feet should be established around these sensitive areas. Where practicable, native vegetation may be an alternative to turf in the buffer areas.

It is noted that identification of sensitive natural resources areas such as wetlands and streams is based on the publicly available National Hydrologic Dataset and the U.S. Fish & Wildlife Service (USFWS) National Wetland Inventory Maps. Field mapping of other wetlands and streams that may exist on the campus was outside the scope of this NMP.

5.0 OTHER TURF MANAGEMENT CONSIDERATIONS

Aeration - Extensive core cultivation/aeration in the late summer to early fall is recommended for the Locust Grove campus. Core aeration is very disruptive to surface smoothness, but it is the best way to relieve the physical effects of soil compaction and increase soil oxygen levels.

Grass Seed Type - Reference the most recent Virginia Cooperative Extension's *Virginia Turfgrass Variety Recommendations* found online at <https://www.sites.ext.vt.edu/newsletter-archive/turfgrass/index.html> when selecting seed mix for over-seeding. The seed type should be suitable to regional environmental conditions.

Iron - Iron applications (particularly foliar applications) may periodically be used for enhanced greening as an alternative to nitrogen. These applications are most beneficial if applied in late spring through summer for cool season grasses and in late summer/fall applications for warm-season grasses. Since iron is a micronutrient, its application levels are very low. The color response is short-lived (typically two to three weeks) because the iron-induced color response in the leaves is removed by mowing.

Returning and Management of Grass Clippings - The recycling of grass clippings on turf should be encouraged as an effective means of recycling nitrogen, phosphorus, and potassium. Where aesthetics allows, all clippings from mowing events should be returned to the turf rather than discharging them onto sidewalks or streets. Clippings should not be blown onto impervious surfaces or surface waters, dumped down stormwater drains, or piled outside where rainwater will leach out the nutrients creating the potential for nutrient loss to the environment

Spreader Equipment Calibration - Spreader equipment calibration is critical to NMP implementation. The landscape contractor should supply equipment calibration records to the Director of Facilities on a routine basis.

6.0 RECORDKEEPING

Proper NMP implementation requires diligent record keeping of fertilizer, lime and herbicide applications, and turfgrass conditions. Important information to retain with the plan includes soil tests reports; spreader settings; calibration results, dates of fertilizer application and rates applied; seeding or renovation; and unusual stresses caused by disease, drought, and pests. This information will also provide the background needed for future plan revisions. NMP Application record keeping forms are included in Appendix B for use for tracking fertilizer, lime, pesticide and herbicides.

7.0 REFERENCES

Nutrient Management Training and Certification Regulations 4VAC50-85 (effective date November 23, 2014)

Virginia Nutrient Management Standards and Criteria (Revised July 2014):

<https://www.dcr.virginia.gov/document/standardsandcriteria.pdf>

Urban Nutrient Management Handbook (August 16, 2019); 430-350: <https://resources.ext.vt.edu/>

A Spreadsheet-Based Soil Test Converter for Turfgrass Professionals and Nutrient Management Planning in Virginia (November 1, 2018); SPES-60P: <https://resources.ext.vt.edu/>

Soil Test Note #1 – Explanation of Soil Tests (December 1, 2018): <https://resources.ext.vt.edu/>

TABLES

Table 1 - Soil Test Summary

Site: Locust Grove Campus - GCC											
Testing Lab: Waypoint Analytical (Formerly A&L Eastern Laboratories)											
Sample Date: 12/6/2024											
Soil Sampling Area ID	Square Feet	Soil pH (SU)	Buffer pH (SU)	P (Mehlich III) ppm	P (Mehlich I) ppm	P (H/M/L)	K (Mehlich III) ppm	K (Mehlich I) ppm	K (H/M/L)	Soil description	Turf Species
LG-1	206,138.99	5.3	6.77	7	0	L-	82	58	M	Dark Brown, Sandy Clay Loam	Cool season, fescue
LG-2	29,386.01	5.7	6.82	8	0	L-	44	31	L+	Dark Brown, Clay Loam	Cool season, fescue
LG-3	149,298.52	5.5	6.85	5	0	L-	28	20	L	Dark Brown, Clay Loam	Cool season, fescue

Notes: SU = Standard Units; ppm = parts per million; P and K ratings are from Virginia Nutrient Management Standards & Criteria.

Table 2 – Nutrient Application Worksheet – Schedule 1**Site:** Locust Grove Campus**Operator:** Germanna Community College**Begins:** 3/01/2025**Expires:** 3/01/2028**Nutrient Management Area:** 1 (LG-1 & LG-2)**Square Feet:** 235,525.00**Landscape Plants:** Cool Season Turf (Fescue)

Annual Nutrient Needs (lbs./1000 ft ²) ¹	Application Month/Day ^{1,2}	Amendment Material Notes	% Slow-Release N	Total N (lbs./1000 ft ²)	Total P ₂ O ₅ (lbs./1000 ft ²)	Total K ₂ O (lbs./1000 ft ²)	Lime Recommendation (lbs./1000 ft ²) ³
3.4 - 3.0 - 1.5	March 4	Lime		0	0	0	35 (LG-1)
	April 4	N	50% or greater	0.7	0	0	
	September 4	N, P ₂ O ₅ , K ₂ O	50% or greater	0.9	1.0	0.50	
	October 4	N, P ₂ O ₅ , K ₂ O	50% or greater	0.9	1.0	0.50	
	November 4	N, P ₂ O ₅ , K ₂ O	50% or greater	0.9	1.0	0.50	
	December 4	Lime		0	0	0	35 (LG-1) & 50 (LG-2)
	Totals:			3.4	3.0	1.5	

Notes:

- Fertilizer recommendations are flexible provided the following conditions are met: a) no more than 0.7 pounds of Water Soluble N per 1000 ft² is applied within a 30-day period; b) no more than 0.9 pounds of Total N (per 1000 ft²) may be applied within a 30-day period; and c) Total annual fertilizer amounts for each nutrient should not exceed the Annual Nutrient Needs listed in column 1.
- The fertilization window is from March 4th until December 8th. The day designations in column 2 are general guidelines. The application day can vary as long as the application intervals are adhered to, and the applications occur within the window dates.
- Lime areas are shown in Figure 4 and soil sampling areas are shown in Figure 2. Two split applications of pelletized agricultural dolomitic limestone at a rate of 35 lbs./1000 ft² for a total of 70 lbs./1000 ft² for soil sampling area LG-1 and one application of 50 lbs./1000 ft² for soil sampling area LG-2. Lime applications are for the first year after sampling only. Liming for the following years should be based on additional soil pH and Buffer pH testing.
- Do not apply inorganic fertilizers on frozen or snow-covered ground, or on denuded areas. Any fertilizer that makes its way onto impervious surfaces should be swept or blown back into pervious turfgrass – covered areas. Do not use fertilizers as ice melt.
- Use a drop spreader for application of inorganic fertilizers on turf areas less than 10 feet wide or on slopes greater than 2%.
- Apply pre and or post emergent herbicides as needed, but do not use fertilizer containing herbicide prior to March 4th. Conditions must be met in Note 1.

Table 3 – Nutrient Application Worksheet – Schedule 2**Site:** Locust Grove Campus**Operator:** Germanna Community College**Begins:** 3/01/2025**Expires:** 3/01/2028**Nutrient Management Area:** 2 (LG-3)**Square Feet:** 149,298.52**Landscape Plants:** Cool Season Turf (Fescue)

Annual Nutrient Needs (lbs./1000 ft ²) ¹	Application Month/Day ^{1,2}	Amendment Material Notes	% Slow-Release N	Total N (lbs./1000 ft ²)	Total P ₂ O ₅ (lbs./1000 ft ²)	Total K ₂ O (lbs./1000 ft ²)	Lime Recommendation (lbs./1000 ft ²) ³
3.4 - 3.0 - 2.5	March 4	Lime		0	0	0	30 (LG-3)
	April 4	N & K ₂ O	50% or greater	0.7	0	0.25	
	September 1	N, P ₂ O ₅ & K ₂ O	50% or greater	0.9	1.0	0.75	
	October 1	N, P ₂ O ₅ & K ₂ O	50% or greater	0.9	1.0	0.75	
	November 1	N, P ₂ O ₅ & K ₂ O	50% or greater	0.9	1.0	0.75	
	December 1	Lime		0	0	0	30 (LG-3)
	Totals:			3.4	3.0	2.5	

Notes:

1. Fertilizer recommendations are flexible provided the following conditions are met: a) no more than 0.7 lbs. of Water Soluble N per 1000 ft² is applied within a 30-day period; b) no more than 0.9 lbs. of Total N (per 1000 ft²) may be applied within a 30-day period; and c) Total annual fertilizer amounts for each nutrient should not exceed the Annual Nutrient Needs listed in column 1.
2. The fertilization window is from March 4th until December 8th. The day designations in column 2 are general guidelines. The application day can vary as long as the application intervals are adhered to, and the applications occur within the window dates.
3. Lime areas are shown in Figure 4 and soil sampling areas are shown in Figure 2. Two split applications of pelletized agricultural dolomitic limestone at a rate of 30 lbs./1000 ft² for a total of 60 lbs./1000 ft² for soil sampling area LG-3. Lime applications are for the first year after sampling only. Liming for the following years should be based on additional soil pH and Buffer pH testing.
4. Do not apply inorganic fertilizers on frozen or snow-covered ground, or on denuded areas. Any fertilizer that makes its way onto impervious surfaces should be swept or blown back into pervious turfgrass – covered areas. Do not use fertilizers as ice melt.
5. Use a drop spreader for application of inorganic fertilizers on turf areas less than 10 feet wide or on slopes greater than 2%.
6. Apply pre and or post emergent herbicides as needed, but do not use fertilizer containing herbicide prior to March 4th. Conditions must be met in Note 1.

FIGURES

Figure 1: Project Location









 <p>Wetland Studies and Solutions, Inc. a DAVEY company</p>	<p> Campus Boundary</p> <p>Prepared by WL 1/17/2025 8:54 AM Sources: VGIN Most Recent Projection: NAD 1983 StatePlane Virginia South FIPS 4502 Feet</p>	<p>FIGURE 1 PROJECT LOCATION GCC - Locust Grove NMP</p> <p>0 1,000 2,000 Feet</p> <p>Locust Grove, Virginia</p>
---	--	---


Figure 2: Soil Sampling Areas







 LG-1

 LG-2

 LG-3

 Campus Boundary

 National Hydrography Dataset (NHD)

Prepared by WL 1/17/2025 9:11 AM
Sources: Google Satellite Imagery
Projection: NAD 1983 StatePlane Virginia South FIPS 4502 Feet

FIGURE 2
SOIL SAMPLING AREAS
GCC - Locust Grove NMP

0

250



500


Feet


Locust Grove, Virginia


Figure 3: Nutrient Management Areas





 NMA-1

 NMA-2

 Campus Boundary


 National Hydrography Dataset (NHD)

FIGURE 3
NUTRIENT MANAGEMENT AREAS
GCC - Locust Grove NMP

0

250

500



Feet

Locust Grove, Virginia




Prepared by WL 1/17/2025 9:51 AM
Sources: Google Satellite Imagery
Projection: NAD 1983 StatePlane Virginia South FIPS 4502 Feet


Figure 4: Liming Areas






Liming Area:

-  50 lbs/1000 ft²
-  60 lbs/1000 ft²*
-  70 lbs/1000 ft²*

 Campus Boundary

 National Hydrography Dataset (NHD)

*See Tables 2 and 3 for split applications

FIGURE 4
LIMING AREA
GCC - Locust Grove NMP

0 250 500 Feet

Locust Grove, Virginia

Prepared by WL 1/17/2025 11:35 AM
Sources: Google Satellite Imagery
Projection: NAD 1983 StatePlane Virginia South FIPS 4502 Feet

APPENDICES

Appendix A: Laboratory Soil Test Results

SOIL ANALYSIS

Client : Wetlands Studies Solutions 1620 Brook Road Richmond VA 23220	Grower : GCC LG GCC Locust Grove Campus 2130 Germanna Highway Locust Grove, VA 22508 PO:	Report No: 24-341-0569 Cust No: 78934 Date Printed: 12/09/2024 Date Received : 12/06/2024 Date Analysis : 12/09/2024 Page : 1 of 6
--	--	---

Lab Number : 11770

Field Id :

Sample Id : LG-1

Test	Results	SOIL TEST RATINGS					Calculated Cation Exchange Capacity
		Very Low	Low	Medium	Optimum	Very High	
Soil pH	5.3						5.2
Buffer pH	6.77						meq/100g
Phosphorus (P)	7 ppm						Calculated Cation Saturation
Potassium (K)	82 ppm						%K 4.0
Calcium (Ca)	479 ppm						%Ca 46.1
Magnesium (Mg)	121 ppm						%Mg 19.4
Sulfur (S)							%H 30.8
Boron (B)							Hmeq 1.6
Copper (Cu)							
Iron (Fe)							
Manganese (Mn)							
Zinc (Zn)							K : Mg Ratio
Sodium (Na)							0.20
Soluble Salts							Ca : Mg Ratio
Organic Matter	4.2 % ENR 127						2.38
Nitrate Nitrogen							

SOIL FERTILITY GUIDELINES

Crop : Lawn

Rec Units: LB/1000 SF

(lbs)	LIME	(tons)	N	P ₂ O ₅	K ₂ O	Mg	S	B	Cu	Mn	Zn	Fe
70			4.0	2.0	2.0	0						
Crop :												Rec Units:

Comment :

Brandi Watson

SOIL ANALYSIS

Client : Wetlands Studies Solutions 1620 Brook Road Richmond VA 23220	Grower : GCC LG GCC Locust Grove Campus 2130 Germanna Highway Locust Grove, VA 22508 PO:	Report No: 24-341-0569 Cust No: 78934 Date Printed: 12/09/2024 Date Received : 12/06/2024 Date Analysis : 12/09/2024 Page : 2 of 6
--	--	---

Lab Number : 11770

Field Id :

Sample Id : LG-1

SUGGESTED FERTILIZATION PROGRAM

First Application		Second Application		Third Application		Fourth Application	
#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer
12	10-20-15	12	10-20-15	8	16-4-8		

Comments:

Lawn

Limestone application is targeted to bring soil pH to 6.5.

- Apply the amount of lime recommended in first page to raise pH
- For existing lawns, lime applications should NOT exceed 50 pounds per 1000 sq. feet per application. Split and apply every 4 to 6 month until the recommended amount is completed.
- The amount of fertilizer recommended on the first page is the total amount needed for the entire growing season. Split into 3-4 applications to keep the lawn green and prevent fertilizer loss. You should not apply more than 0.7 lbs of soluble nitrogen per 1000 square feet in a 30 day period. Or more than 0.9 lbs of nitrogen per 1000 square feet if you are using a slow or controlled release product in a 30 day period. Custom blend is best to meet exactly the requirement, if this is impossible, the above specific fertilizer application is a general guideline, if the specified grades can not be found, replace with fertilizer having similar N:P:K ratio. The best time to apply fertilizer for cool season grass (bluegrass, fescue, ryegrass) is in the Fall when the grass is growing. For Mid-Atlantic region the time is from late August to November. For Northeast region the time is from mid August to October. Fall application should start as soon as the day time high temperature is below 80-85F, apply with the interval of one month. If you start application late in the Fall and do not finish all three applications, repeat the same applications in the Fall of next year. Spring application is recommended when exceptional fertilizer loss due to heavy spring rain leaching and the grasses look pale green. Spring application can start as soon as the grass starts to grow in April. In the case of exceptional warm spring, the application can be made earlier.
- Use calcitic limestone to correct the pH.



SOIL ANALYSIS

Client : Wetlands Studies Solutions 1620 Brook Road Richmond VA 23220	Grower : GCC LG GCC Locust Grove Campus 2130 Germanna Highway Locust Grove, VA 22508 PO:	Report No: 24-341-0569 Cust No: 78934 Date Printed: 12/09/2024 Date Received : 12/06/2024 Date Analysis : 12/09/2024 Page : 3 of 6
--	--	---

Lab Number : 11771

Field Id :

Sample Id : LG-2

Test	Results	SOIL TEST RATINGS					Calculated Cation Exchange Capacity
		Very Low	Low	Medium	Optimum	Very High	
Soil pH	5.7						5.1 meq/100g
Buffer pH	6.82						
Phosphorus (P)	8 ppm						Calculated Cation Saturation
Potassium (K)	44 ppm						%K 2.2
Calcium (Ca)	579 ppm						%Ca 56.8
Magnesium (Mg)	114 ppm						%Mg 18.6
Sulfur (S)							%H 21.6
Boron (B)							Hmeq 1.1
Copper (Cu)							
Iron (Fe)							
Manganese (Mn)							
Zinc (Zn)							K : Mg Ratio
Sodium (Na)							0.10
Soluble Salts							Ca : Mg Ratio
Organic Matter	3.5 % ENR 113						3.05
Nitrate Nitrogen							

SOIL FERTILITY GUIDELINES

Crop : Lawn

Rec Units: LB/1000 SF

(lbs)	LIME	(tons)	N	P ₂ O ₅	K ₂ O	Mg	S	B	Cu	Mn	Zn	Fe
50			4.0	2.0	3.0	0						
Crop :												Rec Units:

Comment :

Brandi Watson

SOIL ANALYSIS

Client : Wetlands Studies Solutions 1620 Brook Road Richmond VA 23220	Grower : GCC LG GCC Locust Grove Campus 2130 Germanna Highway Locust Grove, VA 22508 PO:	Report No: 24-341-0569 Cust No: 78934 Date Printed: 12/09/2024 Date Received : 12/06/2024 Date Analysis : 12/09/2024 Page : 4 of 6
--	--	---

Lab Number : 11771

Field Id :

Sample Id : LG-2

SUGGESTED FERTILIZATION PROGRAM

First Application		Second Application		Third Application		Fourth Application	
#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer
12	10-20-15	12	10-20-15	12	10-0-20		

Comments:

Lawn

Limestone application is targeted to bring soil pH to 6.5.

- Apply the amount of lime recommended in first page to raise pH
- The amount of fertilizer recommended on the first page is the total amount needed for the entire growing season. Split into 3-4 applications to keep the lawn green and prevent fertilizer loss. You should not apply more than 0.7 lbs of soluble nitrogen per 1000 square feet in a 30 day period. Or more than 0.9 lbs of nitrogen per 1000 square feet if you are using a slow or controlled release product in a 30 day period. Custom blend is best to meet exactly the requirement, if this is impossible, the above specific fertilizer application is a general guideline, if the specified grades can not be found, replace with fertilizer having similar N:P:K ratio. The best time to apply fertilizer for cool season grass (bluegrass, fescue, ryegrass) is in the Fall when the grass is growing. For Mid-Atlantic region the time is from late August to November. For Northeast region the time is from mid August to October. Fall application should start as soon as the day time high temperature is below 80-85F, apply with the interval of one month. If you start application late in the Fall and do not finish all three applications, repeat the same applications in the Fall of next year. Spring application is recommended when exceptional fertilizer loss due to heavy spring rain leaching and the grasses look pale green. Spring application can start as soon as the grass starts to grow in April. In the case of exceptional warm spring, the application can be made earlier.
-
- Use calcitic limestone to correct the pH.



SOIL ANALYSIS

Client : Wetlands Studies Solutions 1620 Brook Road Richmond VA 23220	Grower : GCC LG GCC Locust Grove Campus 2130 Germanna Highway Locust Grove, VA 22508 PO:	Report No: 24-341-0569 Cust No: 78934 Date Printed: 12/09/2024 Date Received : 12/06/2024 Date Analysis : 12/09/2024 Page : 5 of 6
--	--	---

Lab Number : 11773

Field Id :

Sample Id : LG-3

Test	Results	SOIL TEST RATINGS					Calculated Cation Exchange Capacity
		Very Low	Low	Medium	Optimum	Very High	
Soil pH	5.5						3.1 meq/100g
Buffer pH	6.85						
Phosphorus (P)	5 ppm						Calculated Cation Saturation
Potassium (K)	28 ppm						%K 2.3
Calcium (Ca)	333 ppm						%Ca 53.7
Magnesium (Mg)	66 ppm						%Mg 17.7
Sulfur (S)							%H 25.8
Boron (B)							Hmeq 0.8
Copper (Cu)							
Iron (Fe)							
Manganese (Mn)							
Zinc (Zn)							K : Mg Ratio
Sodium (Na)							0.17
Soluble Salts							Ca : Mg Ratio
Organic Matter	1.9 % ENR 84						3.03
Nitrate Nitrogen							

SOIL FERTILITY GUIDELINES

Crop : Lawn

Rec Units: LB/1000 SF

(lbs)	LIME	(tons)	N	P ₂ O ₅	K ₂ O	Mg	S	B	Cu	Mn	Zn	Fe
60			4.0	2.0	3.0	0						
Crop :												Rec Units:

Comment :

Brandi Watson

SOIL ANALYSIS

Client : Wetlands Studies Solutions 1620 Brook Road Richmond VA 23220	Grower : GCC LG GCC Locust Grove Campus 2130 Germanna Highway Locust Grove, VA 22508 PO:	Report No: 24-341-0569 Cust No: 78934 Date Printed: 12/09/2024 Date Received : 12/06/2024 Date Analysis : 12/09/2024 Page : 6 of 6
--	--	---

Lab Number : 11773

Field Id :

Sample Id : LG-3

SUGGESTED FERTILIZATION PROGRAM

First Application		Second Application		Third Application		Fourth Application	
#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer	#/1000 Sq. Ft.	Fertilizer
12	10-20-15	12	10-20-15	12	10-0-20		

Comments:

Lawn

Limestone application is targeted to bring soil pH to 6.5.

- Apply the amount of lime recommended in first page to raise pH
- For existing lawns, lime applications should NOT exceed 50 pounds per 1000 sq. feet per application. Split and apply every 4 to 6 month until the recommended amount is completed.
- The amount of fertilizer recommended on the first page is the total amount needed for the entire growing season. Split into 3-4 applications to keep the lawn green and prevent fertilizer loss. You should not apply more than 0.7 lbs of soluble nitrogen per 1000 square feet in a 30 day period. Or more than 0.9 lbs of nitrogen per 1000 square feet if you are using a slow or controlled release product in a 30 day period. Custom blend is best to meet exactly the requirement, if this is impossible, the above specific fertilizer application is a general guideline, if the specified grades can not be found, replace with fertilizer having similar N:P:K ratio. The best time to apply fertilizer for cool season grass (bluegrass, fescue, ryegrass) is in the Fall when the grass is growing. For Mid-Atlantic region the time is from late August to November. For Northeast region the time is from mid August to October. Fall application should start as soon as the day time high temperature is below 80-85F, apply with the interval of one month. If you start application late in the Fall and do not finish all three applications, repeat the same applications in the Fall of next year. Spring application is recommended when exceptional fertilizer loss due to heavy spring rain leaching and the grasses look pale green. Spring application can start as soon as the grass starts to grow in April. In the case of exceptional warm spring, the application can be made earlier.
- Use calcitic limestone to correct the pH.

Brandi Watson

Appendix B: Application Record Forms

Fertilizer Application Records								
Customer Information					Management Area Information			
Name:					Management Area ID:			
Address:					Management Area Size:			
					Target Species:			
					Notes:			
Date	Supervisor/Applicator	Weather Conditions			Fertilizer Analysis	Rate	Amount Fertilizer Used (1000 lbs/AC)	Application Equipment Used
		Temp	Wind Speed	Precip				
When was the last time your fertilizer equipment was calibrated??? For information on calibration see Chapter 10 of the "Urban Nutrient Management Handbook". Available for download at http://pubs.ext.vt.edu/430/430-350/430-350.html								

Herbicide Application Records								
Customer Information					Management Area Information			
Name:					Management Area ID:			
Address:					Management Area Size:			
					Target Species:			
					Notes:			
Date	Supervisor/Applicator	Weather Conditions			Herbicide Analysis	Rate	Amount Herbicide Used	Application Equipment Used
		Temp	Wind Speed	Precip				
When was the last time your fertilizer equipment was calibrated??? For information on calibration see Chapter 10 of the "Urban Nutrient Management Handbook". Available for download at http://pubs.ext.vt.edu/430/430-350/430-350.html								

Lime Application Records								
Customer Information					Management Area Information			
Name:					Management Area ID:			
Address:					Management Area Size:			
					Notes:			
Date	Supervisor/Applicator	Weather Conditions			Lime Analysis	Rate	Amount Lime Used (1000 lbs/AC)	Application Equipment Used
		Temp	Wind Speed	Precip				
When was the last time your fertilizer equipment was calibrated??? For information on calibration see Chapter 10 of the "Urban Nutrient Management Handbook". Available for download at http://pubs.ext.vt.edu/430/430-350/430-350.html								

Pesticide Application Records								
Customer Information					Management Area Information			
Name:					Management Area ID:			
Address:					Management Area Size:			
					Target Species:			
					Notes:			
Date	Supervisor/Applicator	Weather Conditions			Pesticide Analysis	Rate	Amount Pesticide Used	Application Equipment Used
		Temp	Wind Speed	Precip				
When was the last time your fertilizer equipment was calibrated??? For information on calibration see Chapter 10 of the "Urban Nutrient Management Handbook". Available for download at http://pubs.ext.vt.edu/430/430-350/430-350.html								